

# Compactness-like Covering Properties

Petra Staynova

Durham University

November 7, 2013

# Something to think about...

[Look at the board]

One of the main generalisations of compactness is the notion of an H-closed space:

One of the main generalisations of compactness is the notion of an  $H$ -closed space:

## Definition ( $H$ -closed, general)

A topological space  $X$  is said to be  *$H$ -closed* iff it is closed in every Hausdorff space containing it as a subspace.

One of the main generalisations of compactness is the notion of an  $H$ -closed space:

## Definition ( $H$ -closed, general)

A topological space  $X$  is said to be  *$H$ -closed* iff it is closed in every Hausdorff space containing it as a subspace.

A more useful definition is:

## Definition ( $H$ -closed)

A topological space  $X$  is said to be  *$H$ -closed* iff every open cover has a finite subfamily with dense union.

# Countable generalisations

Another generalisation of compactness is the well-known Lindelöf property:

## Definition (Lindelöf)

A topological space  $X$  is called *Lindelöf* iff every open cover has a countable subcover.

# Countable generalisations

Another generalisation of compactness is the well-known Lindelöf property:

## Definition (Lindelöf)

A topological space  $X$  is called *Lindelöf* iff every open cover has a countable subcover.

## Example (Lindelöf spaces)

The following spaces are Lindelöf:

- Any countable topological space;

# Countable generalisations

Another generalisation of compactness is the well-known Lindelöf property:

## Definition (Lindelöf)

A topological space  $X$  is called *Lindelöf* iff every open cover has a countable subcover.

## Example (Lindelöf spaces)

The following spaces are Lindelöf:

- Any countable topological space;
- Any space with the co-countable topology;



# Countable generalisations

Another generalisation of compactness is the well-known Lindelöf property:

## Definition (Lindelöf)

A topological space  $X$  is called *Lindelöf* iff every open cover has a countable subcover.

## Example (Lindelöf spaces)

The following spaces are Lindelöf:

- Any countable topological space;
- Any space with the co-countable topology;
- A countable union of compact spaces;

# Countable generalisations

Another generalisation of compactness is the well-known Lindelöf property:

## Definition (Lindelöf)

A topological space  $X$  is called *Lindelöf* iff every open cover has a countable subcover.

## Example (Lindelöf spaces)

The following spaces are Lindelöf:

- Any countable topological space;
- Any space with the co-countable topology;
- A countable union of compact spaces;
- $\mathbb{R}$ , with the Euclidean topology.

# Countable generalisations

Another generalisation of compactness is the well-known Lindelöf property:

## Definition (Lindelöf)

A topological space  $X$  is called *Lindelöf* iff every open cover has a countable subcover.

## Example (Lindelöf spaces)

The following spaces are Lindelöf:

- Any countable topological space;
- Any space with the co-countable topology;
- A countable union of compact spaces;
- $\mathbb{R}$ , with the Euclidean topology.
- The Sorgenfrey line  $\mathbb{S}$  ( $\mathbb{R}$  with the topology generated by the base  $\mathcal{B} = \{[a, b) : a < b\}$ ).

## Countable generalisations, cont'd

In 1959, Zdenek Frolik introduced a notion that combines the Lindelöf and H-closed properties:

## Countable generalisations, cont'd

In 1959, Zdenek Frolik introduced a notion that combines the Lindelöf and H-closed properties:

### Definition (weakly Lindelöf , [Fro59])

A topological space  $X$  is *weakly Lindelöf* if for every open cover  $\mathcal{U}$  of  $X$  there is a countable subfamily  $\mathcal{U}' \subseteq \mathcal{U}$  such that  $X = \overline{\bigcup \mathcal{U}'}$ .

## Countable generalisations, cont'd

In 1959, Zdenek Frolik introduced a notion that combines the Lindelöf and H-closed properties:

### Definition (weakly Lindelöf , [Fro59])

A topological space  $X$  is *weakly Lindelöf* if for every open cover  $\mathcal{U}$  of  $X$  there is a countable subfamily  $\mathcal{U}' \subseteq \mathcal{U}$  such that  $X = \overline{\bigcup \mathcal{U}'}$ .

Later on, while studying cardinal invariants, Dissanayake and Willard introduced another generalisation of the Lindelöf property, which is stronger than the weakly Lindelöf one:

## Countable generalisations, cont'd

In 1959, Zdenek Frolik introduced a notion that combines the Lindelöf and H-closed properties:

### Definition (weakly Lindelöf , [Fro59])


A topological space  $X$  is *weakly Lindelöf* if for every open cover  $\mathcal{U}$  of  $X$  there is a countable subfamily  $\mathcal{U}' \subseteq \mathcal{U}$  such that  $X = \overline{\bigcup \mathcal{U}'}$ .

Later on, while studying cardinal invariants, Dissanayake and Willard introduced another generalisation of the Lindelöf property, which is stronger than the weakly Lindelöf one:

### Definition (almost Lindelöf , [WD84])


A topological space  $X$  is *almost Lindelöf* iff for every open cover  $\mathcal{U}$  of  $X$  there is a countable subfamily  $\mathcal{U}' \subseteq \mathcal{U}$  such that  $X = \bigcup \{\overline{U} : U \in \mathcal{U}'\}$ .

# Generalisations of compactness: a diagram





# Separation Axioms



## Proposition

- *Every regular Hausdorff  $H$ -closed space is compact.*
- *Every regular Lindelöf space is normal.*

# Properties of the generalisations

## Proposition

- *Every regular Hausdorff  $H$ -closed space is compact.*
- *Every regular Lindelöf space is normal.*

## Proposition

*A regular almost Lindelöf space is Lindelöf.*

# Properties of the generalisations

## Proposition

- *Every regular Hausdorff  $H$ -closed space is compact.*
- *Every regular Lindelöf space is normal.*

## Proposition

*A regular almost Lindelöf space is Lindelöf.*

## Proposition

*A normal weakly Lindelöf space is almost Lindelöf.*

## Proposition

*The continuous image of an almost Lindelöf (weakly Lindelöf) space is almost Lindelöf (weakly Lindelöf).*

# Properties of the generalisations

## Proposition

*The continuous image of an almost Lindelöf (weakly Lindelöf) space is almost Lindelöf (weakly Lindelöf).*

## Proposition

*A clopen subset of an almost Lindelöf (weakly Lindelöf) space is almost Lindelöf (weakly Lindelöf).*

# Properties of the generalisations

## Proposition

*The continuous image of an almost Lindelöf (weakly Lindelöf) space is almost Lindelöf (weakly Lindelöf).*

## Proposition

*A clopen subset of an almost Lindelöf (weakly Lindelöf) space is almost Lindelöf (weakly Lindelöf).*

## Proposition

*If  $X$  is almost Lindelöf (weakly Lindelöf) and  $Y$  is compact, then  $X \times Y$  is almost Lindelöf (weakly Lindelöf).*

Property	Compact	H-closed	Lindelof	Weakly Lindelof
Inherited by closed subspaces?	Yes	No (regularly closed)	Yes	No (regularly closed)



## Definition (quasi-Lindelöf , [Arh79])

We call a space  $X$  *quasi-Lindelöf* if for every closed subset  $Y$  of  $X$  and every collection  $\mathcal{U}$  of open in  $X$  sets such that  $Y \subseteq \bigcup \mathcal{U}$ , there is a countable subfamily  $\mathcal{U}' \subseteq \mathcal{U}$  such that  $Y \subset \overline{\bigcup \mathcal{U}'}$ .

## Definition (quasi-Lindelöf , [Arh79])

We call a space  $X$  *quasi-Lindelöf* if for every closed subset  $Y$  of  $X$  and every collection  $\mathcal{U}$  of open in  $X$  sets such that  $Y \subseteq \bigcup \mathcal{U}$ , there is a countable subfamily  $\mathcal{U}' \subseteq \mathcal{U}$  such that  $Y \subseteq \overline{\bigcup \mathcal{U}'}$ .

## Proposition ([Sta12])

Let  $X$  be a topological space. The following are equivalent:

- 1  $X$  is quasi-Lindelöf.
- 2 Let  $\mathcal{B}$  be a fixed base for  $X$ . Then for any closed subset  $C \subset X$  and any cover  $\mathcal{U}$  of  $C$  with  $\mathcal{U} \subset \mathcal{B}$  there is a countable subfamily  $\mathcal{U}'$  of  $\mathcal{U}$  such that  $C \subseteq \overline{\bigcup \mathcal{U}'}$ .

# weakly Lindelöf not quasi-Lindelöf example

We modify an example from Song and Zhang [SZ10] and use ideas from Mysior [Mys81] to get:

## Example

There exists a Urysohn weakly Lindelöf not quasi-Lindelöf space.

# weakly Lindelöf not quasi-Lindelöf example

We modify an example from Song and Zhang [SZ10] and use ideas from Mysior [Mys81] to get:

## Example

There exists a Urysohn weakly Lindelöf not quasi-Lindelöf space.

(the original example was of a Urysohn almost Lindelöf space which is not Lindelöf)

# Weakly Lindelöf not quasi-Lindelöf example

## Construction

Let  $A = \{(a_\alpha, -1) : \alpha < \omega_1\}$  be an  $\omega_1$ -long sequence in the set  $\{(x, -1) : x \geq 0\} \subseteq \mathbb{R}^2$ . Let  $Y = \{(a_\alpha, n) : \alpha < \omega_1, n \in \omega\}$ . Let  $a = (-1, -1)$ . Finally let  $X = Y \cup A \cup \{a\}$ .

We topologize  $X$  as follows:

# Weakly Lindelöf not quasi-Lindelöf example

## Construction

Let  $A = \{(a_\alpha, -1) : \alpha < \omega_1\}$  be an  $\omega_1$ -long sequence in the set  $\{(x, -1) : x \geq 0\} \subseteq \mathbb{R}^2$ . Let  $Y = \{(a_\alpha, n) : \alpha < \omega_1, n \in \omega\}$ . Let  $a = (-1, -1)$ . Finally let  $X = Y \cup A \cup \{a\}$ .

We topologize  $X$  as follows:

- all points in  $Y$  are isolated;

# Weakly Lindelöf not quasi-Lindelöf example

## Construction

Let  $A = \{(a_\alpha, -1) : \alpha < \omega_1\}$  be an  $\omega_1$ -long sequence in the set  $\{(x, -1) : x \geq 0\} \subseteq \mathbb{R}^2$ . Let  $Y = \{(a_\alpha, n) : \alpha < \omega_1, n \in \omega\}$ . Let  $a = (-1, -1)$ . Finally let  $X = Y \cup A \cup \{a\}$ .

We topologize  $X$  as follows:

- all points in  $Y$  are isolated;
- for  $\alpha < \omega_1$  the basic neighborhoods of  $(a_\alpha, -1)$  will be of the form

$$U_n(a_\alpha, -1) = \{(a_\alpha, -1)\} \cup \{(a_\alpha, m) : m \geq n\} \text{ for } n \in \omega$$

# Weakly Lindelöf not quasi-Lindelöf example

## Construction

Let  $A = \{(a_\alpha, -1) : \alpha < \omega_1\}$  be an  $\omega_1$ -long sequence in the set  $\{(x, -1) : x \geq 0\} \subseteq \mathbb{R}^2$ . Let  $Y = \{(a_\alpha, n) : \alpha < \omega_1, n \in \omega\}$ . Let  $a = (-1, -1)$ . Finally let  $X = Y \cup A \cup \{a\}$ .

We topologize  $X$  as follows:

- all points in  $Y$  are isolated;
- for  $\alpha < \omega_1$  the basic neighborhoods of  $(a_\alpha, -1)$  will be of the form

$$U_n(a_\alpha, -1) = \{(a_\alpha, -1)\} \cup \{(a_\alpha, m) : m \geq n\} \text{ for } n \in \omega$$

- the basic neighborhoods of  $a = (-1, -1)$  are of the form

$$U_\alpha(a) = \{a\} \cup \{(a_\beta, n) : \beta > \alpha, n \in \omega\} \text{ for } \alpha < \omega_1.$$



# Weakly Lindelöf not quasi-Lindelöf example

Let us point out that

## Claim

The subset  $A \subset X$  is closed and discrete in this topology.

# Weakly Lindelöf not quasi-Lindelöf example

Let us point out that

## Claim

The subset  $A \subset X$  is closed and discrete in this topology.

Indeed, for any point  $x \in X$  there is a basic neighborhood  $U(x)$  such that  $A \cap U(x)$  contains at most one point and also that  $X \setminus A = \{a\} \cup Y$  is open (because  $U_\alpha(a) \subset Y \cup \{a\}$ ). Hence  $X$  contains an uncountable closed discrete subset and therefore it cannot be Lindelöf.

# Weakly Lindelöf not quasi-Lindelöf example

Let us point out that

## Claim

The subset  $A \subset X$  is closed and discrete in this topology.

Indeed, for any point  $x \in X$  there is a basic neighborhood  $U(x)$  such that  $A \cap U(x)$  contains at most one point and also that  $X \setminus A = \{a\} \cup Y$  is open (because  $U_\alpha(a) \subset Y \cup \{a\}$ ). Hence  $X$  contains an uncountable closed discrete subset and therefore it cannot be Lindelöf.

Also,

## Note

Note that for any open  $U \ni a$  the set  $X \setminus \bar{U}$  is at most countable.

# Weakly Lindelöf not quasi-Lindelöf example

Let us point out that

## Claim

The subset  $A \subset X$  is closed and discrete in this topology.

Indeed, for any point  $x \in X$  there is a basic neighborhood  $U(x)$  such that  $A \cap U(x)$  contains at most one point and also that  $X \setminus A = \{a\} \cup Y$  is open (because  $U_\alpha(a) \subset Y \cup \{a\}$ ). Hence  $X$  contains an uncountable closed discrete subset and therefore it cannot be Lindelöf.

Also,

## Note

Note that for any open  $U \ni a$  the set  $X \setminus \overline{U}$  is at most countable.

Indeed, for any  $\alpha < \omega_1$ ,  $\overline{U_\alpha(a)} = U_\alpha(a) \cup \{(a_\beta, -1) : \beta > \alpha\}$ . Hence  $X \setminus \overline{U_\alpha(a)}$  is at most countable.

# Weakly Lindelöf not quasi-Lindelöf example

Let us point out that

## Claim

The subset  $A \subset X$  is closed and discrete in this topology.

Indeed, for any point  $x \in X$  there is a basic neighborhood  $U(x)$  such that  $A \cap U(x)$  contains at most one point and also that  $X \setminus A = \{a\} \cup Y$  is open (because  $U_\alpha(a) \subset Y \cup \{a\}$ ). Hence  $X$  contains an uncountable closed discrete subset and therefore it cannot be Lindelöf.

Also,

## Note

Note that for any open  $U \ni a$  the set  $X \setminus \overline{U}$  is at most countable.

Indeed, for any  $\alpha < \omega_1$ ,  $\overline{U_\alpha(a)} = U_\alpha(a) \cup \{(a_\beta, -1) : \beta > \alpha\}$ . Hence  $X \setminus \overline{U_\alpha(a)}$  is at most countable.

It is easily seen that  $X$  is Hausdorff. With a bit more effort, it can also be proven that  $X$  is Urysohn.

# Weakly Lindelöf not quasi-Lindelöf example

## Claim

The space  $X$  is weakly Lindelöf.

# Weakly Lindelöf not quasi-Lindelöf example

## Claim

The space  $X$  is weakly Lindelöf.

Let  $\mathcal{U}$  be an open cover of  $X$ . Then there exists a  $U(a) \in \mathcal{U}$  such that  $a \in U(a)$ . We can find a basic neighborhood  $U_\beta(a) \subset U(a)$ . Then  $\overline{U_\beta(a)} \subset \overline{U(a)}$  and hence  $X \setminus \overline{U(a)}$  will also be at most countable.

# Weakly Lindelöf not quasi-Lindelöf example

## Claim

The space  $X$  is weakly Lindelöf.

Let  $\mathcal{U}$  be an open cover of  $X$ . Then there exists a  $U(a) \in \mathcal{U}$  such that  $a \in U(a)$ . We can find a basic neighborhood  $U_\beta(a) \subset U(a)$ . Then  $\overline{U_\beta(a)} \subset \overline{U(a)}$  and hence  $X \setminus \overline{U(a)}$  will also be at most countable. Hence  $X \setminus \overline{U(a)}$  can be covered by (at most) countably many elements of  $\mathcal{U}$ , say  $\mathcal{U}^*$ . Set  $\mathcal{U}' = \mathcal{U}^* \cup \{U(a)\}$ . Then,  $X \subseteq \bigcup_{U \in \mathcal{U}'} \overline{U} \subseteq \overline{\bigcup_{U \in \mathcal{U}'} U}$ . Therefore,  $X$  is weakly Lindelöf.



# Weakly Lindelöf not quasi-Lindelöf example

## Claim

The space  $X$  is weakly Lindelöf.

Let  $\mathcal{U}$  be an open cover of  $X$ . Then there exists a  $U(a) \in \mathcal{U}$  such that  $a \in U(a)$ . We can find a basic neighborhood  $U_\beta(a) \subset U(a)$ . Then  $\overline{U_\beta(a)} \subset \overline{U(a)}$  and hence  $X \setminus \overline{U(a)}$  will also be at most countable. Hence  $X \setminus \overline{U(a)}$  can be covered by (at most) countably many elements of  $\mathcal{U}$ , say  $\mathcal{U}^*$ . Set  $\mathcal{U}' = \mathcal{U}^* \cup \{U(a)\}$ . Then,  $X \subseteq \bigcup_{U \in \mathcal{U}'} \overline{U} \subseteq \overline{\bigcup_{U \in \mathcal{U}'} U}$ . Therefore,  $X$  is weakly Lindelöf.

## Note

In fact, this shows that  $X$  is even almost Lindelöf.

# Weakly Lindelöf not quasi-Lindelöf example

## Claim

The space  $X$  is not quasi-Lindelöf.

# Weakly Lindelöf not quasi-Lindelöf example

## Claim

The space  $X$  is not quasi-Lindelöf.

Consider the 1-neighborhood of  $a$ :

$$U_1(a) = \{a\} \cup \{(a_\beta, n) : \omega_1 > \beta > 1, n \in \omega\}.$$

We have that  $C = X \setminus U_1(a)$  is closed.

# Weakly Lindelöf not quasi-Lindelöf example

## Claim

The space  $X$  is not quasi-Lindelöf.

Consider the 1-neighborhood of  $a$ :

$$U_1(a) = \{a\} \cup \{(a_\beta, n) : \omega_1 > \beta > 1, n \in \omega\}.$$

We have that  $C = X \setminus U_1(a)$  is closed.

We show the uncountable family of basic open sets

$\mathcal{U} = \{U_0(a_\alpha, -1) : \alpha < \omega_1\}$  forms an open cover of  $C$  which has no countable subcover with dense union.

# Weakly Lindelöf not quasi-Lindelöf example

## Claim

The space  $X$  is not quasi-Lindelöf.

Consider the 1-neighborhood of  $a$ :

$$U_1(a) = \{a\} \cup \{(a_\beta, n) : \omega_1 > \beta > 1, n \in \omega\}.$$

We have that  $C = X \setminus U_1(a)$  is closed.

We show the uncountable family of basic open sets

$\mathcal{U} = \{U_0(a_\alpha, -1) : \alpha < \omega_1\}$  forms an open cover of  $C$  which has no countable subcover with dense union.

Note that the sets  $U_0(a_\alpha, -1)$  are closed and open. Indeed,

$X \setminus U_0(a_\alpha, -1) = \bigcup \{U_0(a_\beta, -1) : \omega_1 > \beta \neq \alpha\} \cup U_{\alpha+1}(a)$ . Hence, if we remove even one of the  $U_0(a_\alpha, -1)$ , the point  $(a_\alpha, -1)$  would remain uncovered. Therefore,  $X$  is not quasi-Lindelöf.

## Another weakly Lindelöf not quasi-Lindelöf example

A shorter example shows that even regularity is not strong enough to make a weakly Lindelöf space quasi-Lindelöf :

# Another weakly Lindelöf not quasi-Lindelöf example

A shorter example shows that even regularity is not strong enough to make a weakly Lindelöf space quasi-Lindelöf :

## Example

The Sorgenfrey plane is weakly Lindelöf but it is not quasi-Lindelöf.

# The quasi-Lindelof property

## Theorem

*Every separable topological space  $X$  is quasi-Lindelöf.*



# The quasi-Lindelof property

## Theorem

*Every separable topological space  $X$  is quasi-Lindelöf.*

## Definition (ccc)

A topological space  $X$  satisfies the *countable chain condition* if every family of non-empty disjoint open subsets of  $X$  is countable.

# The quasi-Lindelof property

## Theorem

*Every separable topological space  $X$  is quasi-Lindelöf.*

## Definition (ccc)

A topological space  $X$  satisfies the *countable chain condition* if every family of non-empty disjoint open subsets of  $X$  is countable.

Arhangel'skii stated without proof that:

## Theorem ([Arh79])

*Every ccc space is weakly-Lindelöf.*

# The quasi-Lindelof property

## Theorem

*Every separable topological space  $X$  is quasi-Lindelöf.*

## Definition (ccc)

A topological space  $X$  satisfies the *countable chain condition* if every family of non-empty disjoint open subsets of  $X$  is countable.

Arhangel'skii stated without proof that:

## Theorem ([Arh79])

*Every ccc space is weakly-Lindelöf.*

In my 3d year project, I proved that:

## Theorem ([Sta11])

*Every ccc space is quasi-Lindelöf.*

## ccc implies quasi-Lindelöf - the proof

Suppose that  $X$  is CCC but not quasi-Lindelöf. Then there exists a closed nonempty set  $F \subset X$  and an uncountable family  $\Gamma = \{U_\alpha : \alpha < \beta\}$  ( $\beta \geq \omega_1$ ) of non-empty open in  $X$  sets such that  $F \subset \bigcup_{\alpha < \beta} U_\alpha$ , but for any countable  $\Gamma' \subset \Gamma$  we have that  $F \setminus \overline{\bigcup \Gamma'} \neq \emptyset$ .

## ccc implies quasi-Lindelöf - the proof

Suppose that  $X$  is CCC but not quasi-Lindelöf. Then there exists a closed nonempty set  $F \subset X$  and an uncountable family  $\Gamma = \{U_\alpha : \alpha < \beta\}$  ( $\beta \geq \omega_1$ ) of non-empty open in  $X$  sets such that  $F \subset \bigcup_{\alpha < \beta} U_\alpha$ , but for any

countable  $\Gamma' \subset \Gamma$  we have that  $F \setminus \overline{\bigcup \Gamma'} \neq \emptyset$ .

We will construct an uncountable collection of nonempty disjoint open in  $X$  sets  $\{V_\gamma : \gamma < \omega_1\}$ , thus contradicting CCC.

## ccc implies quasi-Lindelöf - the proof

Suppose that  $X$  is CCC but not quasi-Lindelöf. Then there exists a closed nonempty set  $F \subset X$  and an uncountable family  $\Gamma = \{U_\alpha : \alpha < \beta\}$  ( $\beta \geq \omega_1$ ) of non-empty open in  $X$  sets such that  $F \subset \bigcup_{\alpha < \beta} U_\alpha$ , but for any

countable  $\Gamma' \subset \Gamma$  we have that  $F \setminus \overline{\bigcup \Gamma'} \neq \emptyset$ .

We will construct an uncountable collection of nonempty disjoint open in  $X$  sets  $\{V_\gamma : \gamma < \omega_1\}$ , thus contradicting CCC.

Let  $V_0 = U_0$ . Then  $F \setminus \overline{U_0} \neq \emptyset$  (and  $X \setminus \overline{U_0} \neq \emptyset$ ). Hence

$$\emptyset \neq F \setminus \overline{U_0} \subset \bigcup \{U_\alpha : \alpha < \beta, \alpha > 0\}$$

and therefore

$$\emptyset \neq F \setminus \overline{U_0} = \bigcup \{U_\alpha \cap (F \setminus \overline{U_0}) : \alpha < \beta, \alpha > 0\}.$$

# ccc implies quasi-Lindelöf - the proof cont'd

Thus we will have  $\alpha_1 \geq 1$  such that  $U_{\alpha_1} \cap (F \setminus \overline{U_0}) \neq \emptyset$  and moreover  $U_{\alpha_1} \cap (X \setminus \overline{U_0}) \neq \emptyset$ .



Thus we will have  $\alpha_1 \geq 1$  such that  $U_{\alpha_1} \cap (F \setminus \overline{U_0}) \neq \emptyset$  and moreover  $U_{\alpha_1} \cap (X \setminus \overline{U_0}) \neq \emptyset$ .

Let  $V_1 = U_{\alpha_1} \cap (X \setminus \overline{U_0})$ . Then  $V_1 \neq \emptyset$ ,  $V_1$  is open in  $X$  and  $V_1 \cap V_0 = \emptyset$ . Again we have

$$\emptyset \neq F \setminus \overline{U_0 \cup U_{\alpha_1}} = \bigcup \{U_\alpha \cap (F \setminus \overline{U_0 \cup U_{\alpha_1}}); \alpha < \beta, \alpha \notin \{0, \alpha_1\}\}.$$

Hence there is  $U_{\alpha_2} \in \Gamma$ ,  $\alpha_1 \notin \{0, \alpha_1\}$  with  $U_{\alpha_2} \cap (F \setminus \overline{U_0 \cup U_{\alpha_1}}) \neq \emptyset$  and moreover  $U_{\alpha_2} \cap (X \setminus \overline{U_0 \cup U_{\alpha_1}}) \neq \emptyset$ .

Thus we will have  $\alpha_1 \geq 1$  such that  $U_{\alpha_1} \cap (F \setminus \overline{U_0}) \neq \emptyset$  and moreover  $U_{\alpha_1} \cap (X \setminus \overline{U_0}) \neq \emptyset$ .

Let  $V_1 = U_{\alpha_1} \cap (X \setminus \overline{U_0})$ . Then  $V_1 \neq \emptyset$ ,  $V_1$  is open in  $X$  and  $V_1 \cap V_0 = \emptyset$ . Again we have

$$\emptyset \neq F \setminus \overline{U_0 \cup U_{\alpha_1}} = \bigcup \{U_\alpha \cap (F \setminus \overline{U_0 \cup U_{\alpha_1}}); \alpha < \beta, \alpha \notin \{0, \alpha_1\}\}.$$

Hence there is  $U_{\alpha_2} \in \Gamma$ ,  $\alpha_1 \notin \{0, \alpha_1\}$  with  $U_{\alpha_2} \cap (F \setminus \overline{U_0 \cup U_{\alpha_1}}) \neq \emptyset$  and moreover  $U_{\alpha_2} \cap (X \setminus \overline{U_0 \cup U_{\alpha_1}}) \neq \emptyset$ .

Define  $V_2 = U_{\alpha_2} \cap (X \setminus \overline{U_0 \cup U_{\alpha_1}})$ . Then  $V_2 \neq \emptyset$ ,  $V_2$  is open in  $X$  and  $V_2$  is disjoint from  $V_0, V_1$ . Let  $\gamma_0 < \omega_1$  and suppose that we have already constructed a family  $\{V_\delta : \delta < \gamma_0\}$  of non-empty, disjoint open in  $X$  sets with  $V_\delta = U_{\alpha_\delta} \cap (X \setminus \overline{\{U_{\alpha_\sigma} : \sigma < \delta\}})$ , where  $\alpha_\delta \notin \{\alpha_\sigma : \sigma < \delta\}$ .

## ccc implies quasi-Lindelöf - the proof cont'd

Since  $\gamma_0$  is a countable ordinal and  $F$  is not weakly Lindelöf in  $X$ , we have that

$$\begin{aligned}\emptyset \neq F \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}} \\ = \bigcup\{U_\alpha \cap (F \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}}) : \alpha < \delta, \alpha \notin \{\alpha_\delta : \delta < \gamma_0\}\}.\end{aligned}$$

## ccc implies quasi-Lindelöf - the proof cont'd

Since  $\gamma_0$  is a countable ordinal and  $F$  is not weakly Lindelöf in  $X$ , we have that

$$\begin{aligned}\emptyset \neq F \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}} \\ = \bigcup\{U_\alpha \cap (F \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}}) : \alpha < \delta, \alpha \notin \{\alpha_\delta : \delta < \gamma_0\}\}.\end{aligned}$$

Hence we can choose  $\alpha_{\gamma_0}$  such that

$$U_{\alpha_{\gamma_0}} \cap (F \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}}) \neq \emptyset$$

and  $\alpha_{\gamma_0} \notin \{\alpha_\delta : \delta < \gamma_0\}$ .

## ccc implies quasi-Lindelöf - the proof cont'd

Since  $\gamma_0$  is a countable ordinal and  $F$  is not weakly Lindelöf in  $X$ , we have that

$$\begin{aligned}\emptyset &\neq F \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}} \\ &= \bigcup\{U_\alpha \cap (F \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}}) : \alpha < \delta, \alpha \notin \{\alpha_\delta : \delta < \gamma_0\}\}.\end{aligned}$$

Hence we can choose  $\alpha_{\gamma_0}$  such that

$$U_{\alpha_{\gamma_0}} \cap (F \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}}) \neq \emptyset$$

and  $\alpha_{\gamma_0} \notin \{\alpha_\delta : \delta < \gamma_0\}$ . Hence moreover

$$U_{\alpha_{\gamma_0}} \cap (X \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}}) \neq \emptyset.$$

Define  $V_{\gamma_0} = U_{\alpha_{\gamma_0}} \cap (X \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}})$ .

Define  $V_{\gamma_0} = U_{\alpha_{\gamma_0}} \cap (X \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}})$ .

Then  $V_{\gamma_0} \neq \emptyset$ ,  $V_{\gamma_0}$  is open in  $X$  and by construction  $V_{\gamma_0} \cap V_\delta = \emptyset$  for every  $\delta < \gamma_0$ .

Define  $V_{\gamma_0} = U_{\alpha_{\gamma_0}} \cap (X \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}})$ .

Then  $V_{\gamma_0} \neq \emptyset$ ,  $V_{\gamma_0}$  is open in  $X$  and by construction  $V_{\gamma_0} \cap V_\delta = \emptyset$  for every  $\delta < \gamma_0$ .

Thus we have constructed a family  $\{V_\gamma : \gamma < \omega_1\}$  of nonempty disjoint open in  $X$  sets, contradicting CCC.



Define  $V_{\gamma_0} = U_{\alpha_{\gamma_0}} \cap (X \setminus \overline{\cup\{U_{\alpha_\delta} : \delta < \gamma_0\}})$ .

Then  $V_{\gamma_0} \neq \emptyset$ ,  $V_{\gamma_0}$  is open in  $X$  and by construction  $V_{\gamma_0} \cap V_\delta = \emptyset$  for every  $\delta < \gamma_0$ .

Thus we have constructed a family  $\{V_\gamma : \gamma < \omega_1\}$  of nonempty disjoint open in  $X$  sets, contradicting CCC.

Hence,  $X$  is quasi-Lindelöf.

# What about the converse?

Though we have that ccc implies quasi-Lindelöf, the converse is false:

## Example

The lexicographic square is quasi-Lindelöf, but is not ccc.

The quasi-Lindelöf property follows from the fact that the lexicographic square is compact, that it is not ccc can be found in [SS96].



# Open questions - preliminaries

The famous Tychonoff's Theorem states that an arbitrary product of compact spaces is compact. However, for Lindelöf spaces, this fails even in the finite case, as exemplified by the Sorgenfrey plane.

## Open questions - preliminaries

The famous Tychonoff's Theorem states that an arbitrary product of compact spaces is compact. However, for Lindelöf spaces, this fails even in the finite case, as exemplified by the Sorgenfrey plane.

In [TAAJ11] Frank Tall introduced the productively Lindelöf property:

### Definition (productively Lindelöf)

A space  $X$  is called *productively Lindelöf* if for every Lindelöf space  $Y$ , the product  $X \times Y$  is Lindelöf.

It is well-known that compact spaces are productively Lindelöf; Tall proved that under certain additional axioms, there exist other productively Lindelöf properties.

Similarly, we can define productively weakly-Lindelöf:

**Definition ((PS) productively weakly Lindelöf)**

A space  $X$  is called *productively weakly Lindelöf* if for every weakly Lindelöf space  $Y$ , the product  $X \times Y$  is weakly Lindelöf.

Similarly, we can define productively weakly-Lindelöf:

## Definition ((PS) productively weakly Lindelöf)

A space  $X$  is called *productively weakly Lindelöf* if for every weakly Lindelöf space  $Y$ , the product  $X \times Y$  is weakly Lindelöf.

## Proposition

*If  $X$  is weakly Lindelöf and  $Y$  is compact, then  $X \times Y$  is weakly Lindelöf.*

This was stated by Song and Zhang in [SZ10], a proof can be found in [Sta11].

We can also define:

**Definition ((PS) productively quasi-Lindelöf )**

A space  $X$  is called *productively quasi-Lindelöf* if for every quasi-Lindelöf space  $Y$ , the product  $X \times Y$  is quasi-Lindelöf.

Thus, it is natural to ask:



# Open questions - continued

We can also define:

## Definition ((PS) productively quasi-Lindelöf )

A space  $X$  is called *productively quasi-Lindelöf* if for every quasi-Lindelöf space  $Y$ , the product  $X \times Y$  is quasi-Lindelöf.

Thus, it is natural to ask:

## Open Question ([Sta12])

Are compact spaces productively quasi-Lindelöf?

This question is interesting even in the partial case:

# Open questions - continued

We can also define:

## Definition ((PS) productively quasi-Lindelöf )

A space  $X$  is called *productively quasi-Lindelöf* if for every quasi-Lindelöf space  $Y$ , the product  $X \times Y$  is quasi-Lindelöf.

Thus, it is natural to ask:

## Open Question ([Sta12])

Are compact spaces productively quasi-Lindelöf?

This question is interesting even in the partial case:

## Open Question ([Sta12])

Is the product of the unit interval  $[0, 1]$  with a quasi-Lindelöf space, quasi-Lindelöf?

## Open questions, cont'd

We know that there is a regular weakly Lindelöf space that is not quasi-Lindelöf, and also that every normal weakly Lindelöf space is quasi-Lindelöf.

We know that there is a regular weakly Lindelöf space that is not quasi-Lindelöf, and also that every normal weakly Lindelöf space is quasi-Lindelöf. Hence, it makes sense to ask:

## Open Question

In completely regular spaces, do the weakly Lindelöf and quasi-Lindelöf properties coincide?

# Bibliography I



A. Arhangel'skii.

A theorem on cardinality.

*Russ. Math. Surv.*, 34(153), 1979.



Z. Frolik.

Generalizations of compact and Lindelöf spaces.

*Czech. Math. J.*, 9, 1959.



A. Mysior.

A regular space which is not completely regular.

*Proceedings of the American Mathematical Society*, 81(4), 1981.



Lynn A. Steen and J. Arthur Seebach.

*Counterexamples in Topology*.

Dover Publications Inc., new edition, 1996.

# Bibliography II



Petra Staynova.

A Comparison of Lindelöf-type Covering Properties of Topological Spaces.

*Rose-Hulman Undergraduate Journal of Mathematics*, 12(2), 2011.



Petra Staynova.

A Note on Quasi-Lindelöf spaces.

*Proceedings of the 41st Spring Conference of the Union of Bulgarian Mathematicians*, 2012.



Y. Song and Y. Zhang.

Some remarks on almost Lindelöf spaces and weakly Lindelöf spaces.

*Matematicheskii Vestnik*, 62, 2010.



Franklin Tall, O. T. Alas, L. F. Aurichi, and L. R. Junqueira.

Non-productively Lindelöf spaces and small cardinals.

*Houston Journal of Mathematics*, 37:1373–1381, 2011.



S. Willard and U. N. B. Dissanayake.

The almost Lindelöf degree.

*Canad. Math. Bull.*, 27(4), 1984.